



Wagner, J. L., Perin, J., Sano Coelho, R., Ting, V. P., Chuck, C. J., & Teixeira Franco, T. (2017). Hydrothermal Conversion of Lipid-Extracted Microalgae Hydrolysate in the Presence of Isopropanol and Steel Furnace Residues. Waste and Biomass Valorization. DOI: 10.1007/s12649-017-9944-7

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1007/s12649-017-9944-7](https://doi.org/10.1007/s12649-017-9944-7)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Springer at <https://link.springer.com/article/10.1007/s12649-017-9944-7>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms.html>

Hydrothermal conversion of lipid-extracted microalgae hydrolysate in the presence of isopropanol and steel furnace residues

Jonathan L. Wagner,^a Julio Perin,^b Renato Sano Coelho,^b Valeska P. Ting,^c

Christopher J. Chuck^{d} and Telma Franco^{*b}*

Supporting Information




Fig. S1: Elemental retention of catalyst material in the solid phase, calculated from SEM-EDX analysis, following HTL reactions; (a) Ladle-furnace residue (LFR), (b) Electric-arc furnace slag (EAFS)

